



Freshwater pests: Parrots feather

Title
Freshwater pests: Parrots feather

Creator
Environmental Reporting, Ministry for the Environment and Statistics New Zealand

Date
2015-10-21

Description
"Freshwater plant and animal pests can have significant negative impacts on ecosystem health by reducing indigenous biodiversity through predation and competition, and destabilising aquatic habitats. Freshwater plant pests can cause economic losses through blocking water intakes for hydroelectricity generation, impeded drainage or irrigation. In addition, pests can affect the suitability for recreational activities. This dataset relates to the "Freshwater pests" measure on the Environmental Indicators, Te taiao Aotearoa website. "

Source
Source: National Institute of Water and Atmospheric Research Method: "Freshwater pest observations, spanning from pre-1980 to 2013, have been mapped. Maps include observation data from three databases: NIWA's Freshwater Fish Database, for fish; NIWA's Freshwater Biodata Information System, for invertebrates and plants; and the Ministry for Primary Industries (MPI)'s Didymo Samples Database, for Didymo. The information in these databases was supplied by a variety of individuals and institutions. Observations were made using a number of methods including line fishing, electrofishing, nets and traps, snorkelling, diving and spotlighting. The accuracy of the data source is of medium quality."

Coverage
-44.8300900079 168.377758011 -35.6158723137 176.828096128

Identifier
<https://data.mfe.govt.nz/layer/52725-freshwater-pests-parrots-feather/>

Language
eng

Subject
New Zealand

Subject
WATER

Subject
FLORA-Exotic

Subject
environment