



Annual average sea surface temperature, 2009

Title

Annual average sea surface temperature, 2009

Creator

Environmental Reporting, Ministry for the Environment and Statistics New Zealand

Date

2015-10-21

Description

"The ocean waters surrounding New Zealand vary in temperature from north to south. They interact with heat and moisture in the atmosphere and affect our weather. Long-term changes and short-term variability in sea-surface temperatures can affect marine processes, habitats, and species. Some species may find it hard to survive in changing environmental conditions. This layer shows annual average sea surface temperature for 2009 as part of the data series for years 1993 to 2013. NIWA's sea-surface temperature archive is derived from the Advanced Very High Resolution Radiometer (AVHRR) satellite data it receives from the National Oceanic and Atmospheric Administration. The archive provides high spatial (approximately 1km) and high temporal (approximately 6-hourly in cloud-free locations) resolution estimates of sea-surface temperatures over the New Zealand region, dating from January 1993. Uddstrom and Oien (1999) and Uddstrom (2003) describe the methods used to derive and validate the data. This dataset relates to the "Annual average sea-surface temperature" measure on the Environmental Indicators, Te taiao Aotearoa website. Geometry: grid Unit: degrees Celsius Further information can be found in: Uddstrom, MJ (2003). Lessons from high-resolution satellite SSTs. Bulletin of the American Meteorological Society, 84(7), 896–897. Uddstrom, MJ, & Oien, NA (1999). On the use of high resolution satellite data to describe the spatial and temporal variability of sea surface temperatures in the New Zealand region. Journal of Geophysical Research (Oceans) 104, chapter 9, 20729–20751. "

Source

Source: National Institute for Water and Atmospheric Research Method: "The NIWA sea surface temperature archive (NSA) is derived from NOAA satellite Advanced Very High Resolution Radiometer (AVHRR) data received by NIWA. It provides high spatial (approximately 1km) and high temporal (approximately 6 hourly in cloud free locations) resolution estimates of sea surface temperatures over the New Zealand region, dating from January 1993. The methods used to derive and validate the NSA are given in Uddstrom and Oien (1999), and Uddstrom (2003). The New Zealand region includes our exclusive economic zone (EEZ), the Chatham Rise, northern subtropical waters, sub Antarctic waters, and the Tasman Sea. It goes from around 30S to 55S, 160E-170W. This data set has been selected as it is representative of the New Zealand region, and the spatial variability of temperature around New Zealand's waters. Globally, oceans have absorbed 30 Units: percent of the warming caused by global greenhouse gas emissions. The accuracy of the data source is of high quality. The data was supplied as a point grid created in Lambert conformal projection and converted to a 0.02 degree raster. "

Coverage

| 55.9144095675 119.03471377 69.6126839709 149.65210644

Identifier

| <https://data.mfe.govt.nz/layer/53101-annual-average-sea-surface-temperature-2009/>

Language

| eng

Subject

| environment

Subject

| New Zealand

Subject

| CLIMATE-AND-WEATHER

Subject

| CLIMATE-AND-WEATHER-Climate-change

Subject

| CLIMATE-AND-WEATHER-Temperature